

## Annex 2

### Exposures that may have put a person at risk of becoming infected with influenza A(H5N1)

Countries and territories where influenza A/H5 viruses have been identified as a cause of illness in human or animal populations since 1 October 2003	Countries and territories where influenza A/H5 viruses have NOT been identified as a cause of illness in human or animal populations since 1 October 2003
<p>During the 7 days before the onset of symptoms, one or more of the following:</p> <ul style="list-style-type: none"> <li>• contact (within 1 metre) with live or dead domestic fowl<sup>1</sup> or wild birds;</li> <li>• exposure to settings where domestic fowl were or had been confined in the previous 6 weeks;</li> <li>• contact (within touching or speaking distance) with a person for whom the diagnosis of influenza A(H5N1) is being considered;</li> <li>• contact (within touching or speaking distance) with a person with an unexplained acute respiratory illness that later resulted in death.</li> </ul>	<p>History of travel, during the 7 days before the onset of symptoms, to a country or territory with reported highly pathogenic avian influenza (HPAI) activity due to influenza A (H5N1) in the animal populations, <b>AND</b> one or more of the following:</p> <ul style="list-style-type: none"> <li>• contact (within 1 metre) with live or dead domestic fowl<sup>1</sup> or wild birds in any setting;</li> <li>• exposure to settings in which domestic fowl were or had been confined in the previous 6 weeks;</li> <li>• contact (within touching or speaking distance) with a confirmed human case of influenza A/H5 infection;</li> <li>• contact (within touching or speaking distance) with a person with an unexplained acute respiratory illness that later resulted in death.</li> </ul>
	<p>During the 7 days before the onset of symptoms, one or more of the following:</p> <ul style="list-style-type: none"> <li>• living in an area in which there are rumours of deaths of domestic fowl;</li> <li>• occupational exposure.<sup>2</sup></li> </ul>
<p>During the 7 days before the onset of symptoms, having worked in a laboratory where there is processing of samples from persons or animals suspected of having HPAI.</p>	

<sup>1</sup> Domestic fowl are birds that are commonly reared for their flesh, eggs, or feathers and are kept in a yard or similar enclosure, including chickens, ducks, geese, turkeys, guinea fowl.

<sup>2</sup> At-risk occupations such as a domestic fowl worker, domestic fowl processing plant worker, domestic fowl culler (catching, bagging, or transporting birds, disposing of dead birds), worker in live animal market, chef working with live or recently killed domestic fowl, dealer or trader in pet birds, health care worker.